

ARTHROSOOTHE™ CREAM

A soothing, topical cream for joint and muscle discomfort



3 oz (90 grams)

Your patients will love this topical cream designed to **alleviate discomfort associated with overused joints and muscles**. It is ideal for arthritis, and should be a staple in everyone's medicine chest.

ArthroSoothe™ Cream contains clinically proven joint health compounds such as Celadrin®, glucosamine, chondroitin and MSM combined with aloe vera gel, emu oil, arnica and other natural compounds known for their healing effects.

Supported by Research-Science First

Celadrin® cream has been clinically shown to reduce inflammation with no side effects. A clinical study involving patients with osteoarthritis conducted at the University of Connecticut showed that topical application of Celadrin® cream produced improvements in the following areas¹:

- Improved knee range of motion
- Unilateral balance
- Ability to walk and sit down
- Ascend/descend stairs
- Ability to rise from sitting

In another study where Celadrin® cream and menthol were tested, one week of treatment was effective in improving functional performance in individuals with arthritis of the knee, wrist and elbow². Research also shows that additional ingredients in ArthroSoothe™ Cream - glucosamine, chondroitin, MSM, aloe vera gel, emu oil and arnica all have anti-inflammatory properties³⁻¹⁰.

How Should ArthroSoothe™ Cream be used?

This rub on cream is a wonderful compliment to our encapsulated ArthroSoothe™ product. Topical application of ArthroSoothe Cream provides quick results with tissue-warming and surface-cooling effects while the capsules work on repairing and lubricating joints.

Recommended use: Apply directly on the forehead, temples or back of neck, muscles, joints or skin anywhere on the body. Avoid contact with eyes and lips. This product is for external use only and may be applied multiple times per day.

Caution: This product should not be used by pregnant women.

Ingredients: Aloe vera gel, Celadrin®, Emu oil, Capric/ caprylic triglycerides, Herbal blend (white willow bark, capsicum, arnica, chamomile, calendula, witch hazel), polysorbate 60, glucosamine, chondroitin, glyceryl monostearate, MSM, glycerin, menthol, cadjuput oil, CMO, carbomer, hyaluronic acid, vitamin E, triethanolamine, balsam of peru oil, phenoxythenol, oil of oregano, peppermint oil, tea tree oil, eucalyptus oil, carnosine.



References:

1. Effect of a cetylated fatty acid topical cream on functional mobility and quality of life of patients with osteoarthritis. Kraemer WJ, Ratamess NA, Anderson JM, Maresh CM, Tiberio DP, Joyce ME, Messinger BN, French DN, Rubin MR, Gómez AL, Volek JS, Hesslink R Jr.; Human Performance Laboratory, School of Medicine, University of Connecticut, Storrs, Connecticut
2. A cetylated fatty acid topical cream with menthol reduces pain and improves functional performance in individuals with arthritis. Kraemer WJ, Ratamess NA, Maresh CM, Anderson JA, Volek JS, Tiberio DP, Joyce ME, Messinger BN, French DN, Sharman MJ, Rubin MR, Gómez AL, Silvestre R, Hesslink RL Jr. *J Strength Cond Res.* 2005 May;19(2):475-80.
3. Glucosamine inhibits IL-1beta-induced NFkappaB activation in human osteoarthritic chondrocytes. Largo R, Alvarez-Soria MA, Díez-Ortego I, Calvo E, Sánchez-Pernaute O, Egido J, Herrero-Beaumont G. *Osteoarthritis Cartilage.* 2003 Apr;11(4):290-8.
4. Glucosamine hydrochloride specifically inhibits COX-2 by preventing COX-2 N-glycosylation and by increasing COX-2 protein turnover in a proteasome-dependent manner. Jang BC, Sung SH, Park JG, Park JW, Bae JH, Shin DH, Park GY, Han SB, Suh SI. *J Biol Chem.* 2007 Sep 21;282(38):27622-32. Epub 2007 Jul 16.
5. Immunomodulatory and anti-inflammatory effects of chondroitin sulphate. du Souich P, García AG, Vergés J, Montell E. *J Cell Mol Med.* 2009 Jun 11
6. Biochemical basis of the pharmacologic action of chondroitin sulfates on the osteoarticular system. Bali JP, Cousse H, Neuzil E. *Semin Arthritis Rheum.* 2001 Aug;31(1):58-68
7. The anti-inflammatory effects of methylsulfonylmethane on lipopolysaccharide-induced inflammatory responses in murine macrophages. Kim YH, Kim DH, Lim H, Baek DY, Shin HK, Kim JK. *Biol Pharm Bull.* 2009 Apr;32(4):651-6.
8. The inner gel component of Aloe vera suppresses bacterial-induced pro-inflammatory cytokines from human immune cells. Habeeb F, Stables G, Bradbury F, Nong S, Cameron P, Plevin R, Ferro VA. *Methods.* 2007 Aug;42(4):388-93.
9. Anti-inflammatory activity of emu oils in rats. Snowden JM, Whitehouse MW. *Inflammopharmacology.* 1997;5(2):127-32.0
10. Screening for anti-inflammatory activity of 12 Arnica (Asteraceae) species assessed by inhibition of NF-kappaB and release of human neutrophil elastase. Ekenäs C, Zebrowska A, Schuler B, Vrede T, Andreasen K, Backlund A, Merfort I, Bohlin L. *Planta Med.* 2008 Dec;74(15):1789-94. Epub 2008 Oct 30.